

## Species *Cohnella colombiensis*

---

### Etymology

[co.lom.bi.en'sis] N.L. fem. adj. *colombiensis*, belonging to Colombia

### Nomenclatural type

[INSDC Nucleotide: CP119317.1](#) <sup>Ts</sup>

### Description

The species is established on the basis of MiGA taxonomic novelty analysis, the taxonomic placement using maximum likelihood trees with 120 bacterial marker genes (bac120) and the type material is the genome MAG\_2441. Genomic metrics include ANI (67.06), AAI (67.21) and dDDH d4 (30.4). The MAG was reconstructed from lignocellulolytic bacterial consortium and encoded for 149 CAZymes, 44 of them involved in lignocellulose degradation. In addition, putative genes for PET and PUR hydrolysis were found within the MAG.

### Classification

*Bacteria* » *Bacillota* » *Bacilli* » *Caryophanales* » *Paenibacillaceae* » *Cohnella* » *Cohnella colombiensis*

### References

Effective publication: Díaz-García et al., 2024 [1]

### Registry URL

<https://seqco.de/i:24096>

## References

1. Díaz-García et al. (2024). Andean soil-derived lignocellulolytic bacterial consortium as a source of novel taxa and putative plastic-active enzymes. *Systematic and Applied Microbiology*. DOI:10.1016/j.syapm.2023.126485