

Sororendozoicomonas aggregata sp. nov.

Submitted by Maire, Justin

Table 1: Complete list of names proposed in the current register list.

Proposed Taxon	Etymology	Description	Parent Taxon	Type	Registry URL
Genus <i>Sororendozoicomonas</i>	[So.ror.en.do.zo.i.co.mo'nas] L. fem. n. <i>soror</i> , sister; L. fem. n. <i>Endozoicomonas</i> , a bacterial genus; L. fem. n. <i>Sororendozoicomonas</i> , Sister taxon of Endozoicomonas genus	<i>Sororendozoicomonas</i> (So.ror.en.do.zo.i.co.mo'nas. L. fem. n. <i>soror</i> , sister; <i>Endozoicomonas</i> , taxonomic name of a bacterial genus; <i>Sororendozoicomonas</i> referring to the close relationship to the bacterial genus <i>Endozoicomonas</i>). The genus <i>Sororendozoicomonas</i> represents a distinct lineage within the Endozoicomonadaceae family, sister to <i>Endozoicomonas</i> , supported by concatenated marker gene phylogenies as well as 16S rRNA gene phylogeny in the present study. <i>Sororendozoicomonas</i> show similar functional and metabolic potential to <i>Endozoicomonas</i> species. The new genus contains two strains isolated from <i>Pocillopora</i> corals (Pac-P3-11-1, from the present study, and Endozoicomonadaceae SCSIO12664) and potentially additional strains isolated from Red Sea <i>Stylophora pistillata</i> (only 16S rRNA gene sequences are available - KC669256.1, KC669230.1, KC669217.1, KC669131.1).	<i>Endozoicomonadaceae</i>	<i>Sororendozoicomonas aggregata</i> ^{Ts}	seqco.de/i:32314

Proposed Taxon	Etymology	Description	Parent Taxon	Type	Registry URL
<p>Species <i>Sororendoziomonas aggregata</i>^{Ts}</p>	<p>[ag.gre.ga'ta] L. part. adj. <i>aggregata</i>, joined together</p>	<p><i>Sororendoziomonas aggregata</i> (ag.gre.ga'ta. L. fem. part. adj. <i>aggregata</i>, joined together, referring to the ability to form aggregates within coral tissues). The new species contains two strains isolated from <i>Pocillopora</i> corals (Pac-P3-11-1, from the present study, and Endozoicomonadaceae SCSIO12664).</p> <p>Type strain: Pac_P3-11-1, isolated from tissues of the coral <i>Pocillopora acuta</i>, collected from Orpheus Island, Australia (18°36'16.26" S, 146°29'24.27" E). Taxonomic assignment based on concatenated marker gene phylogenies as well as 16S rRNA gene phylogeny. Grows on Marine Agar 2216 (Difco) between 23°C and 28°C, forming circular, white, raised, sticky colonies with entire margins.</p>	<p><i>Sororendoziomonas</i></p>	<p>NCBI Assembly: GCA_031528575.1^{Ts}</p>	<p>seqco.de/i:32313</p>